

AD-A140 175


A LABORATORY INVESTIGATION OF THE HIGH-FREQUENCY
FARLEY-BUNEMAN INSTABILITY (U) IOWA UNIV IOWA CITY DEPT
OF PHYSICS AND ASTRONOMY B KUSTOM ET AL. APR 84
U. OF IOWA-84-8 N00014-83-K-0452


1/1

UNCLASSIFIED

F/G 4/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA140175

A LABORATORY INVESTIGATION OF THE HIGH-FREQUENCY
FARLEY-BUNEMAN INSTABILITY

by

B. Kustom, N. D'Angelo, and R. L. Merlino
Department of Physics and Astronomy
The University of Iowa
Iowa City, Iowa 52242

ABSTRACT

A laboratory investigation of the high-frequency Farley-Buneman instability is described. The instability was studied theoretically by Lee et al. [1971] and is predicted to occur in the low E region of the ionosphere when the E/B drift velocity of the electrons relative to the ions is several times C_s , the ion-acoustic speed. In our experiments, an increase of the electric field well above the Lee et al. "threshold" merely enhances the general power level of the fluctuations but does not effect appreciably their spectral shape. The observed frequency spectra fall off in all cases very rapidly with

DTIC
APR 17 1984
A

This document has been approved
for public release and sale; its
distribution is unlimited

DTIC FILE COPY

84 04 16 062

increasing frequency, with a spectral shape of the type $P(f) \propto f^{-3.5}$.

This result has negative implications for a recently proposed mechanism of "anomalous" wave electron heating in the lower E region of the ionosphere.

Accession For

NTIS 68 27

THE

1988

1992

1. *Chlorophyll a* and *Chlorophyll b* contents were determined by spectrophotometry using the method of Lichtenthaler and Whaley (1987).

1990

Dist. .

A1



1. INTRODUCTION

In the ~ 100 to ~ 115 Km height range of the ionosphere $\nu_i > \omega_{ci}$, whereas $\nu_e \ll \omega_{ce}$, where $\nu_{i(e)}$ is the ion (electron) collision frequency with the neutral gas molecules of the atmosphere and $\omega_{ci}(\omega_{ce})$ the ion (electron) gyrofrequency. If an electric field, \underline{E} , is present, normal to the geomagnetic field, \underline{B} , the electrons drift with a velocity $\underline{v_D} = \underline{E} \times \underline{B}/B^2$, while the ions are essentially "held back" by collisions with the neutral gas background. When the velocity of the electrons relative to the ions exceeds the local ion-acoustic speed, C_s , Farley [1963] and Buneman [1963] have shown that longitudinal plasma waves grow, with $K_{\parallel}^2 \ll K_{\perp}^2$ and a phase velocity perpendicular to \underline{B} essentially equal to the electron drift velocity. Here K_{\parallel} and K_{\perp} are the components of the propagation vector, \underline{K} , parallel and perpendicular to \underline{B} , respectively.

The Farley-Buneman instability is responsible for the so-called "Type 1" irregularities observed by radar backscatter in the equatorial and in the auroral electrojet (e.g., Bowles et al. [1963], Balsley [1969], Chesnut [1969]). Rocket measurements of the instability in the polar cap E-region have been reported, e.g., by Olesen et al. [1976] and Bahnsen et al. [1978]. Laboratory experiments on the instability were performed by Saito et al. [1964],

D'Angelo et al. [1974], John and Saxena [1975], Alport et al. [1981], Mikkelsen et al. [1981].

In 1971 Lee, Kennel, and Kindel [Lee et al., 1971] extended the calculations of Buneman [1963] and Farley [1963] to modes of higher frequencies and shorter wavelengths, by retaining the term $K^2 \lambda_{De}^2$ (λ_{De} is the electron Debye length) in the dispersion relation and considering electron drift velocities, relative to the ions, several times larger than the ion thermal speed. They found that, the dispersion relation being now density dependent, there is an electron density threshold for excitation of higher frequency modes. For a given density, the fastest growing mode shifts to higher frequency as v_D increases. For v_D equal to three times the ion thermal speed, the fastest mode has a real frequency $\omega_r \approx 0.7 \sqrt{\omega_{ci} \omega_{ce}}$. This means, typically, wavelengths in the lower E region ionosphere of 10-20 cm, or less, rather than the several meters of the ordinary, low-frequency Farley-Buneman instability.

To the best of our knowledge, only one rather detailed study of this high-frequency instability, by means of rocket-borne instrumentation, has been performed in the high-latitude E region [Kelley and Mozer, 1973]. We shall comment on their results in the last section of this paper.

Related to the high-frequency Farley-Buneman instability is the question of anomalous electron heating in the polar E region, as

reported, e.g., by Schlegel and St. Maurice [1981]. With the incoherent scatter facility at Chatanika, Alaska, they measured electron temperatures of up to 1200°K near the 110-km altitude in the polar E region, in the presence of strong electric fields. They argued that no classical heating process can account for these high temperatures, and suggested that the temperature enhancements are caused by the unstable plasma waves produced in the disturbed polar E region by electric fields $E \geq 50\text{--}60$ mV/m, through the instability discussed by Lee et al. [1971]. An extensive theoretical analysis of the anomalous electron heating is given in a paper by St. Maurice et al. [1981].

In this paper we describe an experiment performed to study in the laboratory the high-frequency Farley-Buneman instability. The experimental set-up is similar to that employed by Alport et al. [1981] in their study of the low-frequency Farley-Buneman instability, except that in the present experiment radial electric fields of up to ~ 5 V/cm (rather than the ~ 0.5 V/cm of Alport et al. [1981]) are utilized, following Lee et al. [1971], to shift the fastest growing mode to frequencies on the order of the lower hybrid frequency.

Section 2 of the paper describes the experimental arrangement, while Section 3 contains the experimental results. In Section 4 these results are discussed, also in relation to the important question of the anomalous electron heating in the lower E region and to other ionospheric observations.

2. EXPERIMENTAL SET-UP

We used, for the present experiment, the same plasma device employed by Alport et al. [1981] in their study of the low-frequency Farley-Buneman instability and of the gradient drift instability. The neutral gas (argon) pressure was still on the order of 10^{-3} torr, while the axial magnetic field was ~ 225 gauss. This insures that the two conditions $\omega_{ci} < v_i$ and $\omega_{ce} \gg v_e$ (i.e., collisional ions and collisionless electrons) are satisfied, as required for the development of both the low-frequency and the high-frequency Farley-Buneman instability. However, with a radial \underline{E} field of ~ 0.5 V/cm, as in the Alport et al. [1981] experiment, the azimuthal drift speed of the electrons, relative to the nearly stationary ions, is only $\sim 2.2 \times 10^5$ cm/s, i.e., only slightly in excess of the ion-acoustic speed, $C_s \sim 2 \times 10^5$ cm/s.

For a study of the high-frequency Farley-Buneman instability radial electric fields several times larger are required. For this reason the device had to be partially modified, to allow us to apply radial electric fields of several V/cm. It was also desirable that the radial field be varied while keeping the plasma density approximately constant.

A schematic diagram of the experimental set-up is shown in Figure 1. A comparison of this figure with Figure 4 of Alport et al.

[1981] will make clear which modifications to the original device were needed to obtain the larger radial electric field. We have added a cylindrical aluminum can, 30 cm in diameter, which is electrically connected to anode rings A_2 and A_3 , and an additional set of filaments (AP, or annular plasma, radial filament structure) mounted on anode ring A_2 . The anode end plate (EP) and ring A_4 are still connected to the vacuum chamber, which is grounded. Plasma and primary electrons from the discharge chamber (right side) stream through the aperture in anode ring A_4 , thus producing a plasma (with a diameter determined mainly by the aperture in A_4) which is terminated in the main chamber on the (grounded) end plate attached to A_1 .

Typically, the main discharge, or center plasma (CP), is operated with a background argon pressure $p \approx 10^{-3}$ torr, a discharge current $I_d^{CP} \approx 1 - 4$ A, a discharge voltage $V_d^{CP} \approx 50$ V, and a magnetic field $B = 225$ gauss in the center of the main chamber. The axial variation of the magnetic field is $\sim 15\%$ over the entire length of the plasma column. The plasma density in the center plasma is, generally, $1 - 5 \times 10^{10} \text{ cm}^{-3}$. The annular plasma is produced by a discharge between the AP filaments and anode rings A_2 , A_3 , and the 30 cm diameter aluminum can. This discharge is operated, typically, with $I_d^{AP} \approx 0.01 \text{ A} - 0.03 \text{ A}$ and $V_d^{AP} \approx 50 \text{ V}$. Thus, only a very tenuous plasma is independently produced in the outer, annular discharge region. The potential of the annular plasma is controlled by means of the outer anode bias V_A . All power supplies for producing and biasing the annular plasma are independent of those for the core plasma.

The operation of our device in its present form is somewhat analogous to that of a standard double-plasma (DP) device. In a DP device [Taylor et al., 1972], a "driver" plasma and a "target" plasma are produced in a common vacuum chamber and are separated by a negatively biased grid which largely prevents the electrons of one plasma from intermixing with those of the other. In our present set-up, a role similar to that of the grid of a DP device is played by the axial magnetic field, which reduces the mobility of all plasma particles and of the primary ionizing electrons. However, a substantial (≤ 1 A) radial current flows generally between the annular plasma and the central plasma.

A radial (inward) electric field is produced when the AP anode structure (A_2 , A_3 , and can) is biased to some positive potential V_A , generally less than ~ 20 V. With V_A applied, the space potential of the annular plasma rises to a value $\geq V_A$. With A_2 and A_4 grounded, the central plasma potential increases somewhat with increasing V_A , but only by a small fraction of V_A . Typical results for the radial profiles of Langmuir probe electron saturation current, I_s , and space potential, V_{sp} , across the central plasma and a large portion of the annular plasma, are shown in Figure 2 ($I_d^{CP} = 1.75$ A, $I_d^{AP} = .025$ A, $V_d^{AP} = V_d^{CP} = 50$ V, $V_A = 15$ V). In the region $0 \text{ cm} < R < +4 \text{ cm}$, the potential profile provides an average radial electric field of ~ 2 V/cm.

At different values of V_A we obtain similar radial potential profiles, with a monotonic increase of the radial electric field with

increasing V_A . This is illustrated in Figure 3, where the difference in space potential, ΔV_{sp} , between $R = 5$ cm and $R = 0$ cm is plotted vs. V_A . Figure 3 was constructed by averaging, for any given V_A , the ΔV_{sp} 's obtained under several somewhat different CP and AP discharge conditions. With our present set-up, evidently we do not observe the saturation of the radial field at $E_r \approx 0.5$ V/cm reported by Alport et al. [1981] (see their Figure 7).

Experiments were performed with an A_4 electrode opening of either 2 cm or 4 cm, thereby producing plasma columns of different diameters. On some occasions, a metal cylinder, of either 2 cm or 4 cm in diameter and 4.5 cm in length, was also attached to A_4 , as shown in Figure 1. The terminating plate connected to A_1 was either 7 cm or 11 cm in diameter. Several combinations of these various electrodes were used in the experiments to be described in Section 3, to ensure that the results obtained were largely independent of any particular electrode configuration. We also used Langmuir probes of different sizes, from a minimum diameter of ~ 0.15 cm to a maximum diameter of ~ 0.5 cm.

Possible effects of the Langmuir probe cable capacitance on the spectral shape of the observed fluctuations were repeatedly checked by either varying the length of the cable or by adding in parallel with the cable a capacitance comparable to that of the cable itself. It was concluded that, in the frequency range investigated, cable capacitance had a negligible effect on the spectral shape of the fluctuations.

3. EXPERIMENTAL RESULTS

From the linear theory of Lee et al. [1971] and under the general experimental conditions described in Section 2, we expected that the frequency spectra of electrostatic fluctuations measured for $E/B \approx$ several times C_s would each exhibit a broad peak at $f \approx (0.5 - 0.7)f_{LH}$ (f_{LH} = lower hybrid frequency), namely around a frequency $f \sim 1500$ KHz.

Figure 4 shows the measured (power) spectra of the potential fluctuations between ~ 100 KHz and ~ 1.8 MHz for $p \approx 1 \times 10^{-3}$ torr, $B \approx 225$ gauss and $V_A = 0, 5, 10,$ and 16 volt. The hole diameter of the A_4 electrode in this case was 2 cm, and the spectra were measured at a fixed radial position of ~ 3 cm where E_r is greatest. Evidently, for all four values of V_A , the power spectra fall very rapidly with increasing frequency, in the same general manner reported by, e.g., D'Angelo et al. [1975] and Alport et al. [1981]. Apparently, in the range of V_A 's investigated, the effect of increasing V_A and, thereby, the radial electric field, is mainly to enhance the general power level of the fluctuations, without appreciably affecting the spectral shape. The same results are obtained when the neutral gas pressure is increased to $\sim 3 \times 10^{-3}$ torr.

In the lower portions of the frequency range, the spectra appear to be characterized by a spectral form of the type $P(f) \propto f^{-3.5}$, in general agreement with the results reported for the standard, low-frequency Farley-Buneman instability, by D'Angelo et al. [1975], John and Saxena [1975], and Alport et al. [1981].

A change in slope appears, for each of the three spectra of Figure 4, at the higher frequencies, the spectra becoming somewhat flatter there than at the lower frequencies. For example, at $V_A = 16$ volt, the spectral change occurs around $f \approx 1$ MHz. However, we feel that this effect is, probably, not due to the onset of the Lee et al.'s instability since, if this were the case, it would be hard to understand why the power at the low frequencies should be ~ 20 dB or so larger than at high frequencies. The general shape of the power spectra shown in Figure 4 was consistently observed, in the present experiment, in all spectra measured under conditions we regard as suitable to the excitation of the high-frequency instability of Lee et al. (1971). These spectra do not seem to be a peculiarity of one special experimental arrangement. Not only were such spectra measured with the various combinations of electrode A_4 hole size and end plate EP size; they were also observed (although in an experiment not designed specifically to test the Lee et al. [1971] calculations) by John and Saxena [1975], in a plasma produced by an RF discharge, and with radial electric fields such that $E/B \approx$ several times C_S .

In Figure 5 we show a set of measurements (A_4 hole size of 4 cm, $p \approx 10^{-3}$ torr, $B \approx 225$ gauss and $V_A = 8$ V) of radial profiles of (a) space potential, V_{sp} , (b) electron saturation current, I_s , to a Langmuir probe, (c) electron temperature, T_e , measured by the probe, and (d) power in the potential fluctuations of frequencies $f = 200, 400, 1000$, and 1400 KHz. From Figure 5 we notice the following:

(1) the largest fluctuation amplitude is observed, at all four frequencies, in the annular spatial region where the radial electric field is largest, and (2) although the electron temperature at the outer edges of the plasma column is as low as $\sim 0.4 - 0.5$ eV (consistent with rapid cooling of the electrons by collisions with neutral gas molecules at pressures $p \approx 10^{-3}$ torr), the electron temperature within the column is in the order of $1.4 - 2$ eV and exhibits four well defined peaks. These peaks in T_e do not occur at the same radial positions where E_r or the fluctuation amplitudes, $\tilde{\phi}$, are largest. Instead, the minima of T_e at $R = \pm 3$ cm are observed at or near the locations where both E_r and $\tilde{\phi}$ are largest. The most straightforward explanation of this effect seems to be that the electron temperature enhancements observed in Figure 5 are not due to electron wave heating nor to resistive heating by the radial Pedersen currents, which must predominantly flow radially inward wherever the radial E field is largest; but, rather, to currents flowing along B, just inside and just outside the annulus of largest E_r , to complete the current paths.

This view is strengthened by similar measurements, performed under somewhat different experimental conditions, and shown in Figure 6. The four T_e peaks are still evident here, although with different magnitudes; and the minima of T_e at $R = \pm 3$ cm again occur at the locations of largest E_r and largest fluctuation amplitudes.

The results presented in this section will be discussed in Section 4, together with the relevant information available from ionospheric observations of the high-frequency Farley-Buneman instability and of the electron heating by large E fields in the lower E region.

4. DISCUSSION AND CONCLUSIONS

As pointed out in Section 3, a shift of γ_{\max} (the maximum growth rate of the Farley-Buneman instability) to higher and higher frequencies as E_r/B is continuously increased to several times C_S , is not apparent from our experimental results. This shift of γ_{\max} is predicted by the linear theory of Lee et al. [1971] and we should have expected it to give rise, in the non-linear evolution of the instability, to spectra dominated by a broad peak slightly below f_{LH} . It seems to us quite surprising, if the Lee et al. [1971] instability is indeed excited as expected, that the instability should give rise in the observed turbulent spectra to no recognizable signature or clue of its presence at the linear stage.

What is the reason for this puzzling result? We believe that the size of the Langmuir probes used to detect the fluctuations and the cables connecting our probes to the spectrum analyzer were appropriate for the expected wavelengths and frequencies. We tried different experimental arrangements all with the same results as far as the spectral shape was concerned. Also, as already noted in Sections 2 and 3, we used different combinations of electrodes A_4 and A_1 to make sure that the observed spectra were not a peculiarity of one particular device configuration.

In addition. John and Saxena [1975], working with an entirely different device and radial E fields large enough to give $E/B \sim$ several times C_s , found also spectra very similar to ours. This strengthens our experimental results, adding to our confidence that they are not device-dependent.

It could be argued that, for some obscure reason, the cylindrical plasma geometry of both our experiment and that of John and Saxena's is not suitable to the excitation of the high-frequency instability. This argument, however, would seem very specious indeed, since we have previously reported, under similar geometrical configuration, the excitation of the low-frequency, long-wavelength Farley-Buneman instability. It would appear that, if at all, a cylindrical geometry of some given radius R for the plasma, would be more prejudicial to the excitation of wavelengths $\gtrsim R$ than of wavelengths $\lesssim R$. This argument becomes even more cogent when possible end effects are considered.

Since we do not know of any obvious flaw in the linear calculations of Lee et al. [1971], we must somehow come to the conclusion that, irrespective of the conditions of linear excitation of the Farley-Buneman instability (namely γ_{\max} largest at low frequency or at high frequency) the turbulent spectrum evolves generally in such a way as to contain most of the energy at the longest wavelengths allowed by the size of the system, and that a $P(K) \propto K^{-3.5}$ spectral shape is, somehow, an almost "universal" shape.

This type of argument would be in line with the ideas prevalent on two-dimensional magnetohydrodynamic turbulence, as presented, e.g., in Knorr [1974].

Is this (tentative) conclusion warranted by what we know about the excitation of the high-frequency instability in the high-latitude ionospheric E region? Here the situation seems also quite puzzling. Olesen et al. [1976] and Bahnson et al. [1978] reported observations, by rocket-borne instruments, of the Farley-Buneman instability in the polar cap E-region. The measured DC electric field was ~ 70 -80 mV/m, large enough to provide electron drift velocities of several times the ion-acoustic speed. The high-frequency Farley-Buneman instability should, thus, have been easily excited in their case. They observed in the ELF region (18 Hz to 180 Hz) very regular wave forms (see, e.g., Figure 4 of Bahnson et al. [1978]), i.e., "a nearly sinusoidal wave form, with a frequency close to 100 Hz." This fact already suggests a spectral shape of the fluctuations in which the wave amplitude drops very rapidly with increasing frequency, since, e.g., a nearly flat spectrum would hardly be reconcilable with "a nearly sinusoidal wave form." The VLF sensor provided a signal amplitude at 7 KHz of $\sim 10^{-7}$ V/m/Hz, at 100 -110 Km altitude. Since the corresponding ELF amplitude is given as ~ 5 mV/m, we may compare the two directly if we assume, conservatively, that the line width around 100 Hz was ~ 50 Hz. This gives for the ELF amplitude a figure of $\sim 7 \times 10^{-4}$ V/m/Hz. Thus the amplitude at ~ 7 KHz is $\sim 10^{-4}$ times smaller than the amplitude at

~ 100 Hz, which indicates a spectral shape with very rapidly decreasing power to the higher frequencies.

On the other hand, Kelley and Mozer [1973], in an earlier paper, have reported the observation of the high-frequency Farley-Buneman instability in the auroral E region, when DC electric fields were present of ~ 70 - 80 mV/m. In particular, a sonagram of the AC electric field showed electrical noise extending from zero frequency all the way to ~ 6.5 KHz, the lower hybrid frequency. AC electric field spectra, in the altitude range 96 Km to 113 Km, were essentially flat from ~ 20 Hz to ~ 3 KHz (see Figure 6 of Kelley and Mozer [1973]). There are, however, certain features in the Kelley and Mozer report which appear rather puzzling. The broad band (50 - 1000 Hz) electric field was detected by two independent probe detectors of different antenna lengths (2.36 m and 4.48 m). The voltages detected by the two instruments were proportional to the antenna length, between altitudes of ~ 95 Km and ~ 120 Km, indicating, as the authors remark, that the detectors responded primarily to waves with wavelengths comparable to or larger than several meters. This conclusion seems to be somewhat at odds with a spectrum which is essentially flat between ~ 20 Hz and ~ 3 KHz. In addition, spectra of $\Delta n/n_0$, the percentage density fluctuation, were also obtained (see Figure 9 of Kelley and Mozer [1973]), and between ~ 20 Hz and ~ 500 Hz these spectra were also rather flat. Now, as Kelley and Mozer [1973] argue, the relation between the electric field fluctuations and the $\Delta n/n_0$ is

$\Delta n/n_0 = ie\Delta E/K(kT_e)$, where K is the wave number of the electrostatic perturbation. For a flat ΔE spectrum, one would have expected an amplitude spectrum of $\Delta n/n_0$ falling off, with increasing K , as $1/K$, which is in apparent contrast with the measured $\Delta n/n_0$ spectra.

Related to the question of the excitation of the Farley-Buneman instability is the "anomalous" heating of the polar E region observed, among others, by Schlegel and St. Maurice [1981]. With the incoherent scatter facility at Chatanika, Alaska, they measured electron temperatures of up to 1200°K near the 110 Km altitude in the polar E region when strong DC electric fields were present. They concluded that no classical heating process could account for these high temperatures, and suggested that the temperature enhancements were probably caused by unstable plasma waves of the Farley-Buneman type. This is an interesting suggestion. The heating mechanism, however, depends on the presence of a sufficiently high power at short wavelengths (tens of cm) in the Farley-Buneman spectrum, since the long wavelength waves must necessarily be ineffective to heat the electrons, as can be seen from the following argument. Schlegel and St. Maurice [1981] estimate that a heat input to the electrons of $1 - 2 \times 10^{-6}$ erg cm⁻³ s⁻¹ is required in order to account for the enhanced electron temperatures. Now, the rate of energy transfer from unstable Farley-Buneman waves to the electrons, Q_w , can be estimated as $Q_w = \epsilon/\tau$, where ϵ is the energy density in the waves and τ a characteristic time for energy transfer. If we write $\epsilon = 1/2(n_0 m_1 v_1^2)$, and $v_1 \sim (\Delta n/n_0)C_s$, where n_0 is the

unperturbed plasma density, $\Delta n/n_0$ the fractional density perturbation due to the waves and C_s the ion-acoustic speed, we find that $\epsilon \sim 2 \times 10^{-10}$ erg cm $^{-3}$, when $n_0 \sim 2 \times 10^5$ cm $^{-3}$, $C_s \sim 300$ m/s and $\Delta n/n_0 \sim 0.2$. This value of ϵ is, in fact, likely to be an overestimate, since $(\Delta n/n_0)$'s which are as large as 20%, even at the longest wavelengths, are uncommon.

The Farley-Buneman wave periods, at the long wavelengths, are on the order of 10^{-2} s, and it seems very unlikely that τ should be any smaller than 10^{-2} s, as otherwise no coherent wave (e.g., Bahnsen et al. [1978]) would ever be observed. In fact, the best estimate of τ is, probably, $\sim 10^{-1}$ s, i.e., about equal to the linear growth time computed by Schmidt and Gary [1973]. We, thus, obtain $Q_w = \epsilon/\tau \sim 2 \times 10^{-9}$ erg cm $^{-3}$ s $^{-1}$, which is smaller than $1 - 2 \times 10^{-6}$ erg cm $^{-3}$ s $^{-1}$ by about three orders of magnitude.

It is clear that, if higher frequency waves are excited, for which τ would presumably be much smaller than the ~ 0.1 s estimated for long wavelengths, the energy transfer time to the electrons would be smaller. However, it can readily be seen that, if the wavenumber power spectra of the electrostatic fluctuations were indeed of the type $P(K) \propto K^{-3.5}$, nothing would be gained by invoking a contribution to electron heating from the short wavelength part of the spectrum. With increasing K , the energy available would decrease much faster than the corresponding increase of $1/\tau$.

It seems to us that a reliable measurement of the power spectrum of the high-frequency Farley-Buneman instability in the lower E region is required, under the condition $E \gtrsim 60 - 70$ mV/m. At present, the two sets of measurements available, namely these of Olesen et al. [1976] and Bahnsen et al. [1978], on the one hand, and of Kelley and Mozer [1973] on the other, appear to give contrasting results. Our present laboratory results as well as those of John and Saxena [1975], are not easily reconcilable with those of Kelley and Mozer [1973].

After this paper was prepared, we learned of a very recent work by Pfaff et al. [1984] on electric field and plasma density measurements in the auroral electrojet. With a DC electric field of 54 mV/m, i.e., large enough for the excitation of the Lee et al. [1971] instability, they measured power spectra of the density and electric field fluctuations in which the power decreased rapidly with increasing wavenumber. They also concluded that there appears to be insufficient power in the short wavelength (tens of cm) fluctuations to make the mechanism of anomalous electron heating [Schlegel and St. Maurice, 1981] operative.

ACKNOWLEDGMENTS

It is our pleasure to acknowledge the skillful technical assistance of Alfred Scheller. This work was supported by the United States Office of Naval Research under contract N00014-83-K-0452 and in part by NASA grant NGL-16-001-043.

REFERENCES

1. Alport, M. J., N. D'Angelo, and H. L. Pecseli, A laboratory experiment on EM backscatter from Farley-Buneman and gradient drift waves, J. Geophys. Res., 86, 7694, 1981.
2. Bahnsen, A., E. Ungstrup, C. G. Fälthammar, V. Fahleson, J. K. Olesen, F. Primdahl, F. Spangslev, and A. Pedersen, Electrostatic waves observed in an unstable polar cap ionosphere, J. Geophys. Res., 83, 5191, 1978.
3. Balsley, B. B., Some characteristics of non two-stream irregularities in the equatorial electrojet, J. Geophys. Res., 74, 2333, 1969.
4. Bowles, K. L., B. B. Balsley, and R. Cohen, Field-aligned E region irregularities identified with acoustic plasma waves, J. Geophys. Res., 68, 2485, 1963.
5. Buneman, O., Excitation of field-aligned sound waves by electron streams, Phys. Rev. Lett., 10, 285, 1963.
6. Chesnut, W. G., Low frequency waves and irregularities in the auroral ionosphere as determined by radar measurements, in Low Frequency Waves and Irregularities in the Ionosphere, edited by N. D'Angelo, D. Reidel, Hingham, Mass., 1969.

7. D'Angelo, N., H. L. Pecseli, and P. I. Petersen, The Farley instability: A laboratory test, J. Geophys. Res., 79, 4747, 1974.
8. D'Angelo, N., H. L. Pecseli, and P. I. Petersen, The K spectrum of ionospheric irregularities, J. Geophys. Res., 80, 1854, 1975.
9. Farley, D. T., A plasma instability resulting in field-aligned irregularities in the ionosphere, J. Geophys. Res., 68, 6083, 1963.
10. John, P. I. and Y. C. Saxena, Observations of the Farley-Buneman instability in a laboratory plasma, Geophys. Res. Lett., 2, 251, 1975.
11. Kelley, M. C. and F. S. Mozer, Electric field and plasma density oscillations due to the high-frequency Hall current two-stream instability in the auroral E region, J. Geophys. Res., 78, 2214, 1973.
12. Knorr, G., Stationary and quasistationary solutions of a guiding-center plasma, Plasma Phys., 16, 423, 1974.
13. Lee, K., C. F. Kennel, and J. H. Kindel, High-frequency Hall current instability, Radio Sci., 6, 209, 1971.
14. Mikkelsen, T., H. L. Pecseli, M. J. Alport, and N. D'Angelo, Experimental investigations of turbulence in magnetized, partially ionized plasmas, in Physics of Auroral Arc Formation, Geophys. Monograph Series, Vol. 25, AGU, 1981.

15. Olesen, J. K., F. Primdahl, F. Spangslev, E. Ungstrup, A. Bahnsen, U. Fablesen, C. G. Fälthammar, and A. Pedersen, Rocket-borne wave, field, and plasma observations in unstable polar cap E-region, Geophys. Res. Lett., 3, 711, 1976.
16. Pfaff, R. F., M. C. Kelley, B. G. Fejer, E. Kudeki, C. W. Carlson, A. Pedersen, and B. Hausler, Electric field and plasma density measurements in the auroral electrojet, J. Geophys. Res., 89, 236, 1984.
17. Saito, S., N. Sato, and Y. Hatta, Low-frequency oscillations in coaxial cylindrical discharge tubes in a magnetic field perpendicular to an electric field, Appl. Phys. Lett., 5, 46, 1964.
18. Schlegel, K. and J. P. St. Maurice, Anomalous heating of the polar E region by unstable plasma waves. 1. Observations, J. Geophys. Res., 86, 1447, 1981.
19. Schmidt, M. J. and S. P. Gary, Density gradients and the Farley-Buneman instability, J. Geophys. Res., 78, 8261, 1973.
20. St. Maurice, J. P., K. Schlegel, and P. M. Banks, Anomalous heating of the polar E region by unstable plasma waves. 2. Theory, J. Geophys. Res., 86, 1453, 1981.
21. Taylor, R. J., K. R. MacKenzie, and H. Ikezi, A large double plasma device for plasma beam and wave studies, Rev. Sci. Instrum., 43, 1675, 1972.

FIGURE CAPTIONS

Figure 1. Schematic diagram of the experimental set-up. (a) topview, (b) central plasma and annular plasma filament structures.

Figure 2. Radial profiles of the space potential, V_{sp} (volt), and of the electron saturation current, I_s (mA). ($I_d^{CP} = 1.75$ A, $I_d^{AP} = 0.025$ A, $V_d^{CP} = V_d^{AP} = 50$ V, $V_A = 15.2$ V, plasma density in the range $10^9 - 10^{10}$ cm $^{-3}$).

Figure 3. The average potential difference, $\overline{\Delta V_{sp}}$ (V), between $R = + 5$ cm and $R = 0$ cm, vs. V_A (V).

Figure 4. Power spectra of the potential fluctuations for $V_A = 0, 5, 10$, and 16 V. ($p = 1 \times 10^{-3}$ torr, $B = 225$ G, $R = + 3$ cm).

Figure 5. Radial profiles of (a) space potential, V_{sp} , (b) electron saturation current, I_s , to a Langmuir probe, (c) electron temperature, T_e , and, (d) power in the potential fluctuations at $f = 200, 400, 1000$, and 1400 KHz. ($I_d^{CP} = 4$ A, $I_d^{AP} = 10$ mA, $V_d^{CP} = V_d^{AP} = 50$ V, $V_A = 8$ V, $p = 1 \times 10^{-3}$ torr, $B = 225$ G). The hole size was 4 cm with a 4.5 cm long metal cylinder connected to A_4 .

Figure 6. As in Fig. 5 with a 4 cm hole size, but without the metal cylinder connected to A_4 .




Figure 1

A-G84-256




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER U of I 84-	2. GOVT ACCESSION NO. AD-A140 175	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A LABORATORY INVESTIGATION OF THE HIGH-FREQUENCY FARLEY-BUNEMAN INSTABILITY		5. TYPE OF REPORT & PERIOD COVERED Progress - April 1984
7. AUTHOR(s) B. Kustom, N. D'Angelo, and R. L. Merlino		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Physics and Astronomy Department The University of Iowa Iowa City, Iowa 52242		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K-0452
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Physics Division Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 1984
		13. NUMBER OF PAGES 25
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted to <u>J. Geophys. Res.</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ionospheric Instabilities Farley-Buneman Instability Type 1, irregularities		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (see reverse side)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

↓
A laboratory investigation of the high-frequency Farley-Buneman instability is described. The instability was studied theoretically by Lee et al. [1971] and is predicted to occur in the low E region of the ionosphere when the E/B drift velocity of the electrons relative to the ions is several times C_s , the ion-acoustic speed. In our experiments, an increase of the electric field well above the Lee et al. "threshold" merely enhances the general power level of the fluctuations but does not effect appreciably their spectral shape. The observed frequency spectra fall-off in all cases very rapidly with increasing frequency, with a spectral shape of the type $P(f) \propto f^{-3.5}$. This result has negative implications for a recently proposed mechanism of "anomalous" wave electron heating in the lower E region of the ionosphere.

↓
also $\propto f^{-3.5}$ power

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

END

FILMED

6-84

DTIC